

Optimizing John Deere See & Spray™

Tristen Avent
Ph.D. Graduate -
Weed Science

Jason Norsworthy
Distinguished Professor -
Weed Science

Tom Barber
Professor - Weed Science

Bob Scott
Professor - Weed Science

Background

John Deere has developed targeted spray technology for post-emergence control of escaped weeds. The See and Spray™ targeting system uses multiple detection settings: lowest, low, medium, high and highest. The chosen setting dictates the confidence of a plant seen by cameras as a weed. The lowest sensitivity excludes small weeds, whereas the highest sensitivity detects most weeds, including small ones.

Weeds vary in size and shape. Arkansas producers have difficulty controlling certain weeds, such as pigweed (*Amaranthus spp.*), grasses, morningglories (*Ipomoea spp.*) and yellow nutsedge (*Cyperus esculentus L.*). Arguably, pigweed is the most concerning weed in corn, cotton and soybean. Given the current state of herbicide resistance across Arkansas, it is imperative that herbicide applications not miss this weed.

Research was conducted across the United States using a See & Spray prototype to determine the efficacy of treating weeds with different sensitivity settings and weed sizes. Research was conducted in Arkansas to determine the impact of preemergence residual strength

and See & Spray™ applications of glufosinate on the weed seedbank in soybean over three years using the lowest and highest sensitivity settings. An earlier fact sheet FSA2225, "[Opportunity Costs to Consider when Adopting See & Spray™ Technology](#)" discusses some of these concepts.

Results

- Larger weeds and higher sensitivity settings increase the probability of detection.
- Morningglories were missed by the spray application the least, followed by grasses, Palmer amaranth, and yellow nutsedge.
- Over a three years, the lowest sensitivity setting allowed more Palmer amaranth escapes from a glufosinate application than the broadcast application and the highest sensitivity setting.

SENSITIVITY	PIGWEEED ESCAPES AFTER 3 YEARS (#/A)
Broadcast	23
Highest	18
Lowest	142


*Arkansas Is
Our Campus*

Visit our website at:
<https://www.uaex.uada.edu>


- The greatest reduction in postemergence herbicide use with See & Spray will be realized if a **STRONG** preemergence herbicide is applied at planting.

	EARLY POSTEMERGENCE	MID POSTEMERGENCE
	Area sprayed	
Strong PRE	72%	88%
Weak PRE	99%	95%


- Use at least the high sensitivity to control pigweed or other highly competitive, resistant-prone weeds.
- Most post-emergence herbicide labels require a reapplication interval. Examples: Glufosinate reapplication is five days in soybean, seven days in corn, and 10 days in cotton. 2,4-D choline reapplication is 12 days in soybean and cotton.
- Palmer amaranth will likely be uncontrollable if missed with the first targeted spray.
- Use the lowest sensitivity when targeting volunteer crops.
- There has been no indication that speed influences the ability to hit weeds, within the range that See & Spray™ applications can occur.




Graph 1. Likelihood of See & Spray™ to treat morningglories effectively.



Graph 2. Likelihood of See & Spray™ to treat grasses effectively.



Graph 3. Likelihood of See & Spray™ to treat pigweed effectively.



Graph 4. Likelihood of See & Spray™ to treat yellow nutsedge effectively.

The graphs show the likelihood of treating morningglories (graph 1), grasses (graph 2),

pigweed (graph 3), and yellow nutsedge (graph 4) with See & Spray™. The different lines represent the different spray sensitivities, and herbicides are provided as a reference for maximum weed sizes recommended by the label. To interpret the graph, pigweed has a 70%, 82% and 95% likelihood of being treated if it is 3 inches tall at the lowest, medium, and highest sensitivities, respectively.

References

- Anonymous (2022) *Basagranherbicide*. UPL NA Inc, King of Prussia, PA, EPA Reg. No. 70506-434.
- Anonymous (2024) *Enlist One herbicide*. Corteva Agriscience, LLC, Indianapolis, IN, EPA Reg. No. 62719-695.
- Anonymous (2023) *Liberty 280SL herbicide*. BASF Corporation, Research Triangle Park, NC. EPA Reg. No. 7969-448.
- Anonymous (2017) *Permit herbicide*. Gowan Company, LLC, Yuma, AZ, EPA Reg. No. 81880-2-10163.
- Anonymous (2020) *Roundup PowerMAX3*. Bayer CropScience, LP, St. Louis, MO, EPA Reg. No. 524-659.

TRISTEN AVENT is a graduate assistant with the University of Arkansas System Division of Agriculture, Fayetteville. **JASON NORSWORTHY** is a distinguished professor in weed science at the University of Arkansas System Division of Agriculture, Fayetteville. **TOM BARBER** and **BOB SCOTT** are both professors of weed science and are located in Lonoke.

FSA2212-PD-6-2025

Pursuant to 7 CFR § 15.3, the University of Arkansas System Division of Agriculture offers all its Extension and Research programs and services (including employment) without regard to race, color, sex, national origin, religion, age, disability, marital or veteran status, genetic information, sexual preference, pregnancy or any other legally protected status, and is an equal opportunity institution.