
Incidence and mechanisms of resistance in *Bemisia tabaci* with special reference to biotype Q

R. Nauen, J. Benting, A. Elbert, D. Rogers, and I. Denholm*

Bayer CropScience AG, Research Insecticides, Monheim, Germany


**Rothamsted Research, Harpenden, UK*

Two important biotypes of *Bemisia tabaci*



Q-type

B-type




Esterase banding pattern to
differentiate biotypes

Bemisia tabaci
(sweetpotato whitefly)

Two important biotypes are B- and Q-types. The most dominant biotype world-wide is B, whereas Q is to a greater or lesser extent restricted to the Mediterranean basin (e.g. Almeria Spain).

Resistance monitoring in Almeria, Spain

Almeria, Spain



Nauen et al. (2002) Pest. Manag. Sci. 58


St. Louis – *Bemisia tabaci*, Q-biotype Symposium – April 3, 2006

Spread of neonicotinoid resistance and Q-biotypes in the Mediterranean basin



Bemisia tabaci

The Mediterranean Basin



505327 (A00849) 11-82

Biochemistry of neonicotinoid resistance



Bemisia tabaci







- ↙ Target site insensitivity
- ↙ Resistance enzyme levels
 - ↙ Esterases
 - ↙ Glutathione S-transferases
 - ↙ Monooxygenases
- ↙ Synergism
- ↙ Metabolism *in vivo*

Biochemistry of imidacloprid resistance in whiteflies (negative systems)






Bemisia tabaci



Monoxygenases confer neonicotinoid resistance in *Bemisia tabaci*



Bemisia tabaci



7-Ethoxycoumarin-O-deethylase activity is a biochemical marker linked to neonicotinoid resistance in *Bemisia tabaci*

The figure displays three scatter plots, each showing the relationship between the LC_{50} (ppm) on the x-axis and AChE inhibition (pmol/30min/mg protein) on the y-axis for a specific insecticide. The x-axis is logarithmic, ranging from 0.1 to 1000 ppm. The y-axis is linear, ranging from 0 to 800 pmol/30min/mg protein. Each plot includes a linear regression line and an R^2 value.

- Imidacloprid:** The chemical structure is shown above the plot. The R^2 value is 0.89. The data points show a strong positive correlation.
- Thiamethoxam:** The chemical structure is shown above the plot. The R^2 value is 0.82. The data points show a strong positive correlation.
- Acetamiprid:** The chemical structure is shown above the plot. The R^2 value is 0.75. The data points show a strong positive correlation.

Isosurface of the Fukui Function - Electrophilic Attack






Figure 1. Isosurfaces of the Fukui function for electrophilic attack on IMI. Three levels of isosurfaces are displayed, 0.005 (green, opaque), 0.001 (yellow, transparent) and 0.0005 (white, transparent)



$$f^{-}(\mathbf{r}) = \rho(\mathbf{r}, N) - \rho(\mathbf{r}, N-1)$$

electrophilic attack Fukui function: electrons are withdrawn from the molecule

Synergism




+ 1000 ppm PB



2d

+ 16 ppm IMI





Inhibitors of monooxygenases

Only one biochemical mechanism of resistance to neonicotinoid insecticides was found in five years of research: **Microsomal monooxygenases**




- 20ml glass vials
- 50,000 dpm coated
- Transfer of 500 adults
- 24h exposure
- Homogenisation
- Centrifugation
- Supernatant TLC



I_{50} , nAChR


1.8 nM



23 nM

Biochemistry of neonicotinoid resistance – molecular analysis of *B. tabaci* P450 genes identified *cyp6A14A* levels linked to neonicotinoid resistance*

Cyp6A 14A-Expression



*Data kindly provided by Dr. Juergen Benting (Bayer CropScience, Monheim, Germany)

Dynamics of Resistance to the Neonicotinoids Acetamiprid and Thiamethoxam in *Bemisia tabaci* (Homoptera: Aleyrodidae)

A. RAMI HOROWITZ,¹ SVETLANA KONTSEDALOV,² AND ISAAC ISHAAYA²




J. Econ. Entomol. 97(6): 2051-2056 (2004)

ABSTRACT The dynamics of resistance in the sweetpotato whitefly, *Bemisia tabaci* (Gennadius), to the neonicotinoids acetamiprid and thiamethoxam was studied extensively in cotton fields in Israel during the cotton-growing seasons 1999–2003. Whitefly strains were collected in early and late seasons mainly in three locations in northern, central, and southern Israel. The whiteflies were assayed under laboratory conditions for susceptibility to neonicotinoids, as part of the Israeli cotton insecticide resistance management strategy. Selections to both acetamiprid and thiamethoxam and cross-resistance between them also were conducted in the laboratory. Although no appreciable resistance to acetamiprid was observed up to 2001, a slight increase of approximately five-fold resistance was detected during 2002 and 2003. However, from 2001 to 2003 thiamethoxam resistance increased >100-fold in the Ayalon Valley and Carmel Coast cotton fields. In cross-resistance assays with both neonicotinoids, the strain that had been selected with thiamethoxam for 12 generations demonstrated almost no cross-resistance to acetamiprid, whereas the acetamiprid-selected strain exhibited high cross-resistance of >500-fold to thiamethoxam.

KEY WORDS *Bemisia tabaci*, neonicotinoids, cross-resistance, acetamiprid, thiamethoxam



Acetamiprid-selected whiteflies are highly cross-resistant to thiamethoxam, but not *vice versa*!



Insecticide cross-resistance in several strains of *Bemisia tabaci*

Strain	Origin	Date	Host plant	Biotype
CRE04-01	Crete	2004	Tomato	Q
BR-JM03	Brazil	2003	Tomato	B
ISR-02	Israel	2002	Ornamentals	B
MEX03-02	Mexico	2003	Tomato	B
TUC-03	USA	2003	Melon	B
→ ESP-00	Spain	2000	Pepper	Q
ALM-1	Spain	1994	Tomato	Q

12 (!) different whitefly insecticides (i.e. covering all relevant chemical classes commercially available) were tested in a leaf-dip bioassay (3-4d) against all strains,

Strain SUD-S (insecticide susceptible)

Compound		Concentration [ppm]						Sum
		200	40	8	1,6	0,32	0,064	
Spiromesifen	Mean	100	100	96	78	45		419
	SEM	0	0	2,67	10	5,5		
Imidacloprid	Mean	100	100	95	90	65	5	455
	SEM	0	0	5	0	25	5	
Thiamethoxam	Mean	100	93	83	53	20		349
	SEM	0	5	5	5	8		
Dinotefuran	Mean	100	100	100	93	60	0	453
	SEM	0	0	0	9,42	5	0	
Pyrimetrozine	Mean	100	100	95	95	60	45	495
	SEM	0	0	5	5	10	5	
Pyriproxyfen	Mean	86	52	47	7			192
	SEM	1,33	7,33	6,5	1,78			
Buprofezin	Mean	100	100	100	71	42	0	413
	SEM	0	0	0	3,11	3,5	0	
Pyridaben	Mean	95	85	35	25			240
	SEM	5	5	5	10			
Deltamethrin	Mean	70	30	25	20			145
	SEM	20	10	5	0			
Monocrotophos	Mean	100	47	18				165
	SEM	0	14,81	6,01				
Endosulfan	Mean	100	98	50				248
	SEM	0	6	25				

* nymphs

Strain ESP-00 (Q-type)

Compound		Concentration [ppm]						Sum
		200	40	8	1,6	0,32	0,064	
Spiromesifen	Mean	95	99	72	57	37	0	360
	SEM	3,33	0,88	2,67	1	8,67	0	
Imidacloprid	Mean	57	27	10	0			94
	SEM	15,56	4,44	0	0			
Thiamethoxam	Mean	87	23	3	0			113
	SEM	6,67	4,44	4,44	0			
Dinotefuran	Mean	100	73	45	27	0		245
	SEM	0	11,11	4,44	4,44	0		
Pyrimetrozine	Mean	53	47	17	0			117
	SEM	22,22	11,11	5	0			
Pyriproxyfen	Mean	73	44	19	7			143
	SEM	1,5	4,5	5	6,5			
Buprofezin	Mean	73	54	47	17			191
	SEM	3,5	2	4,5	9,5			
Pyridaben	Mean	100	63	0				163
	SEM	0	11,1	0				
Deltamethrin	Mean	43	0					43
	SEM	24,44	0					
Monocrotophos	Mean	53	17	7				77
	SEM	4,44	8,89	4,44				
Endosulfan	Mean	100	57	40	17			214
	SEM	0	4,44	6,67	4,44			
Flonicamid	Mean	73	20	0	0			93
	SEM	4,44	4,44	0	0			

* nymphs


Neonicotinoid cross-resistance in *Bemisia tabaci*



IMI = Imidacloprid

TMX = Thiamethoxam


DTF = Dinotefuran



$$y = 14592x - 54,2$$

$$R^2 = 0,9289$$


TMX



$$y = 0,472x + 203,26$$

$$R^2 = 0,79$$

IMI



$$y = 0,6622x + 186,2$$


$$R^2 = 0,7048$$

TMX

● Q-type
○ B-type

Correlated are efficacy indices for the compounds displayed. An efficacy index results from mortality figures scored at different concentrations and subsequently summarized.

Alignment of DFT/BP/SVP/COSMO optimized geometries of neonicotinoids




The alignment was done by minimization of the mutual spatial distance of three pharmacophoric points, namely

- (i) the positively charged C-atom connected to the N-NO₂ and N-CN moiety
- (ii) the nitro/cyano groups themselves
- (iii) the nitrogens of the aromatic rings

Dinotefuran


Pymetrozine is clearly cross-resistant to neonicotinoids in *Bemisia tabaci* as results from leaf-dip bioassays revealed

IMI = Imidacloprid
PYM = Pymetrozine



Correlated are efficacy indices for the compounds displayed. An efficacy index results from mortality figures scored at different concentrations and subsequently summarized.



Efficacy of **spiromesifen** and **endosulfan** against *Bemisia tabaci* is not correlated w/ neonicotinoid resistance as leaf-dip bioassays revealed



Correlated are efficacy indices for the compounds displayed. An efficacy index results from mortality figures scored at different concentrations and subsequently summarized.

IMI = Imidacloprid
SPM = Spiromesifen
EDS = Endosulfan

Spiromesifen* is highly active against neonicotinoid resistant whiteflies and a new chemical tool in IRM



- ➔ Active against whitefly nymphs (all stages)
- ➔ Active against adults at high concentrations
- ➔ Reduction of adult fecundity (fewer eggs)
- ➔ Transovarial effects on egg hatch
- ➔ New IRAC MoA Group 23

* **o·b·e·r·o·n**
SMART PERFORMANCE!

NEW
JUDO



Bayer CropScience

Summary

- There is cross-resistance between all neonicotinoid insecticides in field-collected strains of *B. tabaci*, i.e. selecting with one will impact performance of all others
- Pymetrozine seems to be cross-resistant to neonicotinoids
- Neonicotinoids are most likely detoxified by oxidative metabolism as elevated P450 levels, synergist studies and *in vivo* metabolism experiments revealed
- Neonicotinoid resistance in B- and Q-biotypes is conferred by different cytochrome P450's
- No target site insensitivity yet found
- Spiromesifen is a new chemical option and tool for whitefly resistance management